

p -Adic pseudodifferential operators and p -adic wavelets

S.V.Kozyrev

February 7, 2008

Abstract

We introduce a new wide class of p -adic pseudodifferential operators. We show that the basis of p -adic wavelets is the basis of eigenvectors for the introduced operators.

1 Introduction

In the present paper we introduce the family of pseudodifferential operators, which generate p -adic diffusions. The constructed operators, unlike the Vladimirov operator of p -adic fractional derivation [1], will not be diagonalizable by the Fourier transform. These operators will be diagonal in the basis of p -adic wavelets, introduced in [2]. Therefore we will call these operators non translationally invariant (i.e., non commutative with the operators of shift).

In the present paper we consider only the problem of existence of the mentioned operators. To show that the introduced operators T generate diffusions, one has to show that the corresponding evolutions e^{-tT} exist, conserve the cone of positive functions and the total number of particles. Formally these conditions are satisfied, the rigorous proof will be considered elsewhere.

Ultrametric diffusions were considered in [1], [3], [4], [5], [6]. In [7], [8], [9] ultrametric diffusions were applied for description of the relaxation processes in disordered systems, for instance, in spin glasses and proteins. In the present paper we describe diffusions with more general generators, compared to the considered in the mentioned above literature.

The field of p -adic numbers is the simplest example of an ultrametric space. p -Adic mathematical physics, see [1]–[16], studies the models of mathematical physics with the help of p -adic analysis.

The results of the present paper have the following physical applications.

In the works [7], [14] p -adic analysis was applied to describe the replica symmetry breaking, for discussion of the replica approach see [19]. It was shown that the Parisi matrix describing the replica symmetry breaking (before the $n \rightarrow 0$ limit) is a discrete analogue of a p -adic pseudodifferential operator. In the present paper

we introduce a new family of p -adic pseudodifferential operators. The discrete analogues of the introduced operators will be the new examples of the replica matrices. The application of these replica matrices to disordered models (for which the $n \rightarrow 0$ limit for the introduced operators have to be developed) would be an interesting problem.

Another possible application of the approach of the present paper is the application to random walk on hierarchical landscapes of energy, relaxation phenomena in complex systems and basin to basin kinetics, see [22] – [25], [26].

The structure of the present paper is as follows.

In Section 2 we remind some results from p -adic analysis and describe the basis of p -adic wavelets.

In Section 3 we introduce the family of pseudodifferential operators and prove that these operators are diagonal in the basis of p -adic wavelets.

In Section 4 we discuss in more details some particular examples of pseudodifferential operators and compute the asymptotics of population of unit disk for the corresponding ultrametric diffusion.

In Section 5 we in more details discuss the possible physical applications of the results of the present paper.

2 p -Adic analysis

Remind some notations used in p -adic analysis, see [1] for details.

An ultrametric space is a metric space with the metrics $\|\cdot\|$ satisfying the strong triangle inequality

$$\|xy\| \leq \max(\|xz\|, \|yz\|), \quad \forall z$$

The simplest example of ultrametric space is the field Q_p of p -adic numbers. The field Q_p of p -adic numbers is the completion of the field of rational numbers Q with respect to p -adic norm on Q . This norm is defined in the following way. An arbitrary rational number x can be written in the form $x = p^{\gamma \frac{m}{n}}$ with m and n not divisible by p . The p -adic norm of the rational number $x = p^{\gamma \frac{m}{n}}$ is equal to $|x|_p = p^{-\gamma}$. The strong triangle inequality for p -adic numbers takes the form

$$|x + y|_p \leq \max(|x|_p, |y|_p).$$

p -Adic numbers are in one to one correspondence with the (convergent in p -adic norm) series of the form

$$x = \sum_{\gamma=\gamma_0}^{\infty} x_i p^i, \quad x_i = 0, \dots, p-1 \quad (1)$$

The ring Z_p of p -adic integers contains p -adic numbers of the form

$$x = \sum_{\gamma=0}^{\infty} x_i p^i, \quad x_i = 0, \dots, p-1$$

and the group Q_p/Z_p we identify with the set containing fractions

$$x = \sum_{\gamma=\gamma_0}^{-1} x_i p^i, \quad x_i = 0, \dots, p-1$$

The Vladimirov operator of p -adic fractional derivation is defined as follows [1]

$$D^\alpha f(x) = \frac{1}{\Gamma_p(-\alpha)} \int_{Q_p} \frac{f(x) - f(y)}{|x - y|_p^{1+\alpha}} dy \quad (2)$$

where dy is the Haar measure on Q_p and $\Gamma_p(-\alpha)$ is a constant. This operator is diagonalizable by the p -adic Fourier transform F :

$$D^\alpha f(x) = F^{-1} \circ |k|_p^\alpha \circ F f(x)$$

p -Adic wavelets were introduced in [2], where the following theorem was proven.

Theorem. *The set of functions $\{\psi_{\gamma j n}\}$:*

$$\psi_{\gamma j n}(x) = p^{-\frac{\gamma}{2}} \chi(p^{\gamma-1} j x) \Omega(|p^\gamma x - n|_p), \quad \gamma \in \mathbf{Z}, \quad n \in Q_p/Z_p, \quad j = 1, \dots, p-1 \quad (3)$$

is an orthonormal basis in $L^2(Q_p)$ of eigenvectors of the operator D^α :

$$D^\alpha \psi_{\gamma j n} = p^{\alpha(1-\gamma)} \psi_{\gamma j n} \quad (4)$$

Here $\Omega(x)$ is the indicator of the interval $[0, 1]$, which implies for the characteristic function of p -adic disk the expression

$$\Omega(|x|_p) = \begin{cases} 1, & |x|_p \leq 1, \\ 0, & |x|_p > 1. \end{cases}$$

The function $\chi(x)$ is the complex valued character of the p -adic argument which for x given by (1) is defined as

$$\chi(x) = \exp \left(2\pi i \sum_{\gamma=\gamma_0}^{-1} x_i p^i \right)$$

The theorem above shows that the p -adic wavelets are the basis of eigenvectors for the operator (2). The operator (2), see [1], was considered as the generator of the ultrametric diffusion. In the present paper we introduce a more general class of p -adic pseudodifferential operators. These operators can not be diagonalized by the p -adic Fourier transform, but we will show, that the introduced operators are still diagonalizable by the p -adic wavelet transform (i.e. the p -adic wavelets are eigenvectors of the introduced operators).

p -Adic numbers are related to the Bruhat–Tits trees, see [27]. The Bruhat–Tits tree is the infinite tree (with the infinite number of edges and vertices) where each vertex is connected with exactly $p + 1$ over vertices. p -Adic projective line, which

contains the field of p -adic numbers plus one infinite point, is equivalent to the space of classes of equivalence of semiinfinite paths in the Bruhat–Tits tree. Two (semiinfinite) paths are equivalent if they coincide, starting from some vertex. For instance, the set of semiinfinite paths going through the fixed vertex and through p edges incident to the vertex is homeomorphic to a p -adic disk.

This construction is the analogue of the Poincare model of the Lobachevskii plane, where the Bruhat–Tits tree is the analogue of the Lobachevskii plane, the p -adic (projective) line is the analogue of the absolute, and the (unique) path in the Bruhat–Tits tree connecting x and y in Q_p is the analogue of the geodesic line connecting the points at the absolute.

We introduce the level, or height of the vertex, in the following way. We put in correspondence to every vertex in the Bruhat–Tits tree the p -adic disk, corresponding to the following set of semiinfinite paths in the tree which begin in the vertex. This set contains all the paths going through the p of $p + 1$ edges incident to the vertex (excluding one edge incident to the path going through the infinite point of the absolute). We call the level of the vertex the $\log_p$ of the radius of this disk.

For two points x, y in Q_p there exists the unique path in the Bruhat–Tits tree connecting these two points. This path contains the unique point $A(x, y)$ with the highest level.

3 Construction of the pseudodifferential operators

In [28], [29] examples of ultrametric diffusions on a discrete set were considered. In [1], [3], [4]–[6], [8]–[9] examples of p -adic diffusions were investigated.

In the present section we introduce a family of p -adic pseudodifferential operators, which generate ultrametric diffusions on the field Q_p of p -adic numbers. The existence of corresponding diffusions (possibility to take the exponent of the generator) will not be discussed in the present paper. The introduced pseudodifferential operators will be the generalizations of the Vladimirov operator.

Consider the operator

$$Tf(x) = \int T(x, y)(f(x) - f(y))dy \quad (5)$$

Here we take x, y lying in the field Q_p of p -adic numbers. We consider the operator T as the operator in the space $L^2(Q_p)$ of complex valued square integrable functions of p -adic argument.

In the following we will introduce some conditions for the kernel $T(x, y)$. Operator of the form (5) satisfying these conditions we call the generator of the ultrametric diffusion. We show that under these conditions the operator (5) will be diagonal in the basis of p -adic wavelets, introduced in [2]. This means that the assumption of ultrametricity of the diffusion makes the problem of investigation of the diffusion exactly solvable.

Definition 1. We call the transition probability (and the corresponding diffusion with the generator given by (5)) ultrametric, if the transition probability $T(x, y)$ depends only on the highest point $A(x, y)$ lying at the path between x and y in the Bruhat–Tits tree.

Notice that the transition probability $T(x, y)$ is symmetric with respect to x, y and satisfies the following property:

Lemma 2. The function $T(x, y)$ is symmetric and positive with respect to y , and, for fixed x , is locally constant with respect to y outside any vicinity of x . Moreover, for fixed x , the following condition is satisfied:

$$T(x, y) = \text{const} , \quad \text{if} \quad |x - y|_p = \text{const} \quad (6)$$

Vice versa, when the condition (6) is satisfied, the function $T(x, y)$ satisfies the conditions of Definition 1.

It is easy to see that the Vladimirov operator and similar operators with translationally invariant kernels $T(x, y)$ (which depend only on the difference $x - y$ and therefore, under the corresponding conditions of integrability of the kernel, can be diagonalized by the Fourier transform), satisfy the conditions of lemma 2. In the present paper we introduce the examples of the generators with non translationally invariant kernels.

The following theorem gives the general form of the kernel for the operator (5).

Theorem 3. The function of the form

$$T(x, y) = \sum_{\gamma n} T^{(\gamma n)} \delta_{1, |p^\gamma x - p^\gamma y|_p} \Omega(|p^\gamma x - n|_p) \quad (7)$$

where $T^{(\gamma n)} \geq 0$, $\gamma \in \mathbb{Z}$, $n \in Q_p/Z_p$ satisfies the conditions of lemma 2.

Moreover, an arbitrary function satisfying (6) can be represented in the form (7).

Proof Positivity of $T(x, y)$ is obvious.

Let us prove the symmetricity of $T(x, y)$. We have

$$T(x, y) - T(y, x) = \sum_{\gamma n} T^{(\gamma n)} \delta_{1, |p^\gamma x - p^\gamma y|_p} (\Omega(|p^\gamma x - n|_p) - \Omega(|p^\gamma y - n|_p)) \quad (8)$$

Let us prove that this expression is equal identically to zero. Consider the case when x is such that the following characteristic function is non-zero:

$$\Omega(|p^\gamma x - n|_p) = 1$$

This implies

$$|x - p^{-\gamma} n|_p \leq p^{-\gamma} \quad (9)$$

If

$$\delta_{1, |p^\gamma x - p^\gamma y|_p} \neq 0$$

then

$$|x - y|_p = p^\gamma \quad (10)$$

Formulas (9), (10) and ultrametricity imply that

$$\Omega(|p^\gamma y - n|_p) = 1$$

Therefore the corresponding terms in (8) cancel. This proves that $T(x, y)$ given by (7) is symmetric.

Prove that $T(x, y)$ given by (7) satisfies (6). Fix $x \in Q_p$. Then for y lying at the sphere with the center in x and the radius p^γ we have

$$\delta_{1, |p^\gamma x - p^\gamma y|_p} = 1$$

and

$$\Omega(|p^\gamma x - n|_p)$$

will be equal to 0 or 1 depending on whether $p^{-\gamma}n$ lies in the disk

$$\{y : |x - y|_p \leq p^\gamma\}$$

or no. Therefore $T(x, y)$ will be a constant on the considered sphere and (6) will be satisfied.

This proves that $T(x, y)$ satisfying the conditions of the present lemma will satisfy lemma 2.

Vice versa, it is easy to see that the kernel (7) for x, y lying in the disk with the center in $p^{-\gamma}n$ and the radius p^γ , and satisfying $|x - y|_p = p^\gamma$, takes the value $T(\gamma n)$.

Since all the space $x, y \in Q_p \times Q_p$ is the disjoint union of such a subsets, therefore, taking an arbitrary positive $T(\gamma n)$ we are able to construct an arbitrary kernel satisfying (6). This finishes the proof of the theorem.

In the next theorem we compute the eigenvalues of the generator of ultrametric diffusion.

Theorem 4. *Let the kernel (6) satisfy the condition of convergence of all the integrals in (12) for any γ, n . Then the operator (5) is a well defined operator in $L^2(Q_p)$ with the dense domain and the p -adic wavelets $\psi_{\gamma jn}$ are eigenvectors for the ultrametric diffusion:*

$$T\psi_{\gamma jn} = \lambda_{\gamma n}\psi_{\gamma jn} \quad (11)$$

with the eigenvalues

$$\lambda_{\gamma n} = \int_{|n - p^\gamma y|_p > 1} T(p^{-\gamma}n, y) dy + p^\gamma T(p^{-\gamma}n, p^{-\gamma}(n + 1)) \quad (12)$$

Proof To prove the present theorem we use Lemma 2. Consider the wavelet $\psi_{\gamma jn}$. Then

$$T\psi_{\gamma jn}(x) = \int T(x, y) (\psi_{\gamma jn}(x) - \psi_{\gamma jn}(y)) dy$$

Consider the following cases.

1) Let $|p^\gamma x - n|_p > 1$. Then Lemma 2 implies

$$T\psi_{\gamma jn}(x) = \int T(x, y)\psi_{\gamma jn}(x)dy - T(x, n) \int \psi_{\gamma jn}(y)dy = 0$$

2) Let $|p^\gamma x - n|_p \leq 1$. Then again by Lemma 2

$$\begin{aligned} T\psi_{\gamma jn}(x) &= \left(\int_{|p^\gamma x - p^\gamma y|_p > 1} + \int_{|p^\gamma x - p^\gamma y|_p = 1} + \int_{|p^\gamma x - p^\gamma y|_p < 1} \right) T(x, y)(\psi_{\gamma jn}(x) - \psi_{\gamma jn}(y))dy = \\ &= \left(\int_{|p^\gamma x - p^\gamma y|_p > 1} + \int_{|p^\gamma x - p^\gamma y|_p = 1} \right) T(x, y)(\psi_{\gamma jn}(x) - \psi_{\gamma jn}(y))dy = \\ &= \psi_{\gamma jn}(x) \int_{|p^\gamma x - p^\gamma y|_p > 1} T(x, y)dy + \int_{|p^\gamma x - p^\gamma y|_p = 1} T(x, y)(\psi_{\gamma jn}(x) - \psi_{\gamma jn}(y))dy = \\ &= \psi_{\gamma jn}(x) \int_{|n - p^\gamma y|_p > 1} T(p^{-\gamma}n, y)dy + \sum_{l=0}^{p-1} \int_{|z|_p < p^\gamma} T(x, x + p^{-\gamma}l + z)(\psi_{\gamma jn}(x) - \psi_{\gamma jn}(x + p^{-\gamma}l + z))dz = \\ &= \psi_{\gamma jn}(x) \int_{|n - p^\gamma y|_p > 1} T(p^{-\gamma}n, y)dy + p^{\gamma-1} \sum_{l=1}^{p-1} T(x, x + p^{-\gamma}l)(\psi_{\gamma jn}(x) - \psi_{\gamma jn}(x + p^{-\gamma}l)) \end{aligned}$$

The last equality follows from Lemma 2 and the local constance of $\psi_{\gamma jn}$.

Using the identity

$$\psi_{\gamma jn}(x + p^{-\gamma}l) = p^{-\frac{\gamma}{2}} \chi(p^{\gamma-1}j(x + p^{-\gamma}l)) \Omega(|p^\gamma(x + p^{-\gamma}l) - n|_p) = \chi(p^{-1}jl) \psi_{\gamma jn}(x)$$

and Lemma 2, we obtain

$$T\psi_{\gamma jn}(x) = \psi_{\gamma jn}(x) \left(\int_{|n - p^\gamma y|_p > 1} T(p^{-\gamma}n, y)dy + p^{\gamma-1} T(p^{-\gamma}n, p^{-\gamma}(n+1)) \sum_{l=1}^{p-1} (1 - \chi(p^{-1}jl)) \right)$$

Since

$$\sum_{l=1}^{p-1} (1 - \chi(p^{-1}jl)) = p$$

we obtain

$$T\psi_{\gamma jn}(x) = \psi_{\gamma jn}(x) \left(\int_{|n - p^\gamma y|_p > 1} T(p^{-\gamma}n, y)dy + p^\gamma T(p^{-\gamma}n, p^{-\gamma}(n+1)) \right)$$

which gives (12).

Therefore the operator T is well defined on the basis in $L^2(Q_p)$ that finishes the proof of the theorem. Note that convergence of all the integrals used in the proof is guaranteed by convergence of (12).

The next corollary gives a simple representation for the eigenvalues of the generator with the kernel (7).

Corollary 5. *Let the following series converge for any n :*

$$\sum_{\gamma=0}^{\infty} p^{\gamma} T^{(\gamma n)} < \infty \quad (13)$$

Then the operator (5) corresponding to the kernel (7) is well defined in $L^2(Q_p)$, diagonal in the basis of p -adic wavelets and have the eigenvalues

$$\lambda_{\gamma n} = p^{\gamma} T^{(\gamma n)} + (1 - p^{-1}) \sum_{\gamma'=\gamma+1}^{\infty} p^{\gamma'} T^{(\gamma', p^{\gamma'-\gamma} n)} \quad (14)$$

Note that condition of convergence of the series (13) is equivalent to convergence of the integral (12).

Proof Substituting (7) into (12) we get

$$\begin{aligned} \lambda_{\gamma n} &= \int_{|n-p^{\gamma}y|_p > 1} \sum_{\gamma' n'} T^{(\gamma' n')} \delta_{1, |p^{\gamma'} p^{-\gamma} n - p^{\gamma'} y|_p} \Omega(|p^{\gamma'} p^{-\gamma} n - n'|_p) dy + \\ &\quad + p^{\gamma} \sum_{\gamma' n'} T^{(\gamma' n')} \delta_{1, |p^{\gamma'} p^{-\gamma} n - p^{\gamma'} p^{-\gamma} (n+1)|_p} \Omega(|p^{\gamma'} p^{-\gamma} n - n'|_p) = \\ &= \sum_{\gamma' n'} T^{(\gamma' n')} p^{\gamma'} (1 - p^{-1}) \theta(\gamma' - \gamma) \Omega(|p^{\gamma'} p^{-\gamma} n - n'|_p) + p^{\gamma} \sum_{n'} T^{(\gamma n')} \Omega(|n - n'|_p) = \\ &= p^{\gamma} T^{(\gamma n)} + (1 - p^{-1}) \sum_{\gamma'=\gamma+1}^{\infty} p^{\gamma'} T^{(\gamma', p^{\gamma'-\gamma} n)} \end{aligned}$$

Here we use the properties

$$\int_{|n-p^{\gamma}y|_p > 1} \delta_{1, |p^{\gamma'} p^{-\gamma} n - p^{\gamma'} y|_p} dy = p^{\gamma'} (1 - p^{-1}) \theta(\gamma' - \gamma), \quad \theta(\gamma) = 0, \gamma \leq 0, \theta(\gamma) = 1, \gamma > 0$$

$$\begin{aligned} \delta_{1, |p^{\gamma'} p^{-\gamma} n - p^{\gamma'} p^{-\gamma} (n+1)|_p} &= \delta_{\gamma \gamma'} \\ \Omega(|m - n|_p) &= \delta_{mn}, \quad m, n \in Q_p / Z_p \end{aligned}$$

This finishes the proof of the corollary.

Remark. This corollary implies the recurrent formulas

$$\begin{aligned} \lambda_{\gamma-1, p^{-1}n} - \lambda_{\gamma n} &= p^{\gamma-1} T^{(\gamma-1, p^{-1}n)} - p^{\gamma-1} T^{(\gamma n)} \\ T^{(\gamma n)} &= T^{(\gamma-1, p^{-1}n)} + p^{1-\gamma} (\lambda_{\gamma n} - \lambda_{\gamma-1, p^{-1}n}) \end{aligned} \quad (15)$$

For instance the sequence $\{\lambda_{\gamma n}\}$ for fixed n is non-negative and non-increasing with the increase of γ .

4 Examples of the pseudodifferential operators and properties of the diffusions

Calculate particular examples of the kernels of the kind (7).

Example 1. Let us consider the translationally invariant case. In this case

$$T^{(\gamma n)} = T^{(\gamma)}$$

and the summation over n in (7) gives

$$\sum_n \Omega(|p^\gamma x - n|_p) = 1$$

Denoting

$$T^{(\gamma)} = f(p^\gamma)$$

we obtain for the kernel

$$T(x, y) = \sum_\gamma T^{(\gamma)} \delta_{1, |p^\gamma x - p^\gamma y|_p} = f(|x - y|_p)$$

Therefore the choice

$$T^{(\gamma)} = p^{-\gamma(1+\alpha)} \tag{16}$$

corresponds to the Vladimirov operator, when the kernel is given by

$$T(x, y) = \frac{1}{|x - y|_p^{1+\alpha}}$$

Formula (14) in the considered case (16) takes the form

$$\begin{aligned} \lambda_{\gamma n} &= p^\gamma T^{(\gamma)} + (1 - p^{-1}) \sum_{\gamma'=\gamma+1}^{\infty} p^{\gamma'} T^{(\gamma')} = \\ &= p^\gamma p^{-\gamma(1+\alpha)} + (1 - p^{-1}) \sum_{\gamma'=\gamma+1}^{\infty} p^{\gamma'} p^{-\gamma'(1+\alpha)} = p^{-\gamma\alpha} \frac{1 - p^{-\alpha-1}}{1 - p^{-\alpha}} \end{aligned} \tag{17}$$

Example 2. Consider the case when

$$T^{(\gamma n)} = f(p^\gamma) g(|p^{-\gamma} n - n_0|_p) \tag{18}$$

This implies for the kernel

$$\begin{aligned} T(x, y) &= \sum_\gamma f(p^\gamma) \delta_{1, |p^\gamma x - p^\gamma y|_p} \sum_n g(|p^{-\gamma} n - n_0|_p) \Omega(|p^\gamma x - n|_p) = \\ &= f(|x - y|_p) \sum_\gamma \delta_{p^\gamma, |x-y|_p} \sum_n g(|p^{-\gamma} n - n_0|_p) \Omega(|p^\gamma x - n|_p) \end{aligned}$$

One can check that the series

$$\sum_{n \in Q_p/Z_p} g(|n - n_0|_p) \Omega(|x - n|_p)$$

equals to $g(|x - n_0|_p)$ for $|x - n_0|_p > 1$ and to $g(0)$ for $|x - n_0|_p \leq 1$:

$$\sum_{n \in Q_p/Z_p} g(|n - n_0|_p) \Omega(|x - n|_p) = g(|x - n_0|_p) (1 - \Omega(|x - n_0|_p)) + g(0) \Omega(|x - n_0|_p)$$

This implies

$$\begin{aligned} & \sum_{n \in Q_p/Z_p} g(|p^{-\gamma}n - n_0|_p) \Omega(|p^\gamma x - n|_p) = \\ & = g(|x - n_0|_p) (1 - \Omega(|p^\gamma x - n_0|_p)) + g(0) \Omega(|p^\gamma x - n_0|_p) \end{aligned}$$

For the kernel this will give

$$T(x, y) = f(|x - y|_p) (g(|x - n_0|_p) (1 - \Omega(|x - y|_p x - n_0|_p)) + g(0) \Omega(|x - y|_p x - n_0|_p)) \quad (19)$$

Corollary 5 and (18) imply for the eigenvalues corresponding to diffusion operator with the kernel (19) the following:

$$\lambda_{\gamma n} = p^\gamma f(p^\gamma) g(|p^{-\gamma}n - n_0|_p) + (1 - p^{-1}) g(|p^{-\gamma}n - n_0|_p) \sum_{\gamma'=\gamma+1}^{\infty} p^{\gamma'} f(p^{\gamma'})$$

In the present section we investigate the relaxation of the indicator function of the unit disk:

$$\phi = \Omega(|x|_p)$$

for the ultrametric diffusion with the kernel (7).

The indicator function has the following decomposition over p -adic wavelets, see [2]

$$\langle \Omega(|x|_p), \psi_{\gamma j n} \rangle = p^{-\frac{\gamma}{2}} \theta(\gamma) \delta_{n0}, \quad \theta(\gamma) = 0, \gamma \leq 0, \quad \theta(\gamma) = 1, \gamma \geq 1$$

$$\Omega(|x|_p) = \sum_{j=1}^{p-1} \sum_{\gamma=1}^{\infty} p^{-\frac{\gamma}{2}} \psi_{\gamma j 0}(x)$$

Therefore the operator T acts on $\Omega(|x|_p)$ as follows

$$T\Omega(|x|_p) = \sum_{j=1}^{p-1} \sum_{\gamma=1}^{\infty} p^{-\frac{\gamma}{2}} \lambda_{\gamma 0} \psi_{\gamma j 0}$$

By (14) for the survival probability of the unit disk with the center in zero we obtain

$$\begin{aligned} S(t) &= \langle \Omega(|x|_p), e^{-tT} \Omega(|x|_p) \rangle = (p-1) \sum_{\gamma=1}^{\infty} p^{-\gamma} e^{-t\lambda_{\gamma 0}} = \\ &= (p-1) \sum_{\gamma=1}^{\infty} p^{-\gamma} \exp\left(-t \left(p^\gamma T^{(\gamma 0)} + (1 - p^{-1}) \sum_{\gamma'=\gamma+1}^{\infty} p^{\gamma'} T^{(\gamma' 0)} \right)\right) \end{aligned} \quad (20)$$

Analogously

$$\begin{aligned}\langle \Omega(|p^\gamma x - n|_p), \psi_{\gamma'jn'} \rangle &= \langle \Omega(|p^\gamma x - n|_p), p^{-\frac{\gamma'}{2}} \chi(p^{\gamma'-1} jx) \Omega(|p^{\gamma'} x - n'|_p) \rangle = \\ &= p^{-\frac{\gamma'}{2}} \theta(\gamma' - \gamma) \chi(p^{\gamma'-\gamma-1} jn) p^\gamma \Omega(|p^{\gamma'-\gamma} n - n'|_p)\end{aligned}$$

which implies

$$\Omega(|p^\gamma x - n|_p) = p^\gamma \sum_{j=1}^{p-1} \sum_{\gamma'=\gamma+1}^{\infty} p^{-\frac{\gamma'}{2}} \chi(p^{\gamma'-\gamma-1} jn) \psi_{\gamma'jp^{\gamma'-\gamma}n}(x) \quad (21)$$

where we use the formula

$$\Omega(|p^{\gamma'-\gamma} n - n'|_p) = \delta_{p^{\gamma'-\gamma}n, n'}$$

Formula (21) allows to compute the analogues of (20) for displaced disks.

Note that $p^{\gamma'-\gamma}n = 0$ in $\mathbb{Q}_p/\mathbb{Z}_p$ when γ' is large enough. Thus the series over the wavelets in (21) contains only the finite sum of the wavelets with the third index not equal to zero. Therefore the asymptotic dependence on time of

$$\langle \Omega(|p^\gamma x - n|_p), e^{-tT} \Omega(|p^{\gamma'} x - n'|_p) \rangle$$

coincides with the asymptotics of (20).

5 Conclusion

In the present paper we construct the new family of p -adic pseudodifferential operators T and show that these operators are diagonal in the basis of p -adic wavelets. This allows to compute analytically the corresponding evolutions e^{-tT} .

The approach of the present paper may be applied to the problem of investigation of the diffusion on hierarchical energy landscapes. The concept of hierarchical energy landscape attracts a lot of interest in connection with investigation of complex systems, in particular, glasses, clusters, and proteins [17]-[26]. Discuss this concept using the language of basis to basis kinetics [26]. In the framework of this concept a complex system is assumed to have a large number of metastable configurations, corresponding to local minima of energy. The local minima are clustered in hierarchically nested basins of minima, namely, each large basin consists of smaller basins, each of these consisting of even smaller ones, and so on. In the other words, the hierarchy of basins possesses ultrametric geometry. Finally, the basins of minima are separated from one another by a hierarchically arranged set of barriers, i.e., high barriers separate large basins, and smaller basins within each larger one are separated by lower barriers.

The concept of hierarchical energy landscape may be considered as a kind of approximation of a complex energy landscape by a hierarchical energy landscape.

The dynamics of the system with hierarchical energy landscape it is natural to describe by the models of ultrametric diffusion.

It is commonly recognized that relevant analytical tools should be developed for applying the concept of hierarchical energy landscape to the description of relaxation phenomena in complex systems [22] – [25], [26] (the concept of “basin-to-basin kinetics”).

The present paper is devoted to the development of these analytical tools. We show, that for some wide class of ultrametric diffusions the corresponding generators (p -adic pseudodifferential operators) may be diagonalized in the basis of p -adic wavelets, introduced in [2]. This means that the random walk on the corresponding energy landscape may be investigated analytically.

Acknowledgements

The author would like to thank V.S.Vladimirov, V.A.Avetisov, I.V. Volovich, A.Yu.Khrennikov, A.H.Bikulov and V.A.Osipov for discussions and valuable comments. This work has been partly supported by INTAS YSF 2002–160 F2, CRDF (grant UM1–2421–KV–02), The Russian Foundation for Basic Research (projects 02–01–01084), and by the grant for the support of scientific schools NSh 1542.2003.1.

References

- [1] V.S.Vladimirov, I.V.Volovich, Ye.I.Zelenov, *p-Adic analysis and mathematical physics*, Moscow, Nauka, 1994; V.S.Vladimirov, I.V.Volovich, Ye.I.Zelenov, *p-Adic analysis and mathematical physics*, Singapore, World Scientific, 1994
- [2] S.V.Kozyrev, Wavelet analysis as a p -adic spectral analysis, *Izvestia Akademii Nauk Seria Math.*, 2002, **66**, N2, <http://xxx.lanl.gov/abs/math-ph/0012019>
- [3] S.Albeverio, W.Karwowski, A random walk on p -adic numbers, pp.61–74 in “Stochastic Process–Physics and Geometry II”, Proc. Locarno (1991), edited by S.Albeverio, U.Cattaneo, D.Merlini, Singapore, World Scientific, 1995
- [4] A.N.Kochubei, On p -adic Green functions, *Teoreticheskaya i Matematicheskaya Fizika*, 1993, **96** N1, pp.123–138
- [5] A.N.Kochubei, Parabolic equations over the field of p -adic numbers, *Izvestia Akademii Nauk Seria Math.*, 1991, **55** N6, pp.1312–1330; A.N.Kochubei, *Math.USSR Izv.*, 1992, **39**, pp.1263–1280
- [6] A.N.Kochubei, *Pseudodifferential equations and stochastics over non-archimedean fields*, New York, Marcel Dekker, 2001
- [7] V.A.Avetisov, A.H.Bikulov, S.V.Kozyrev, Application of p -adic analysis to models of spontaneous breaking of the replica symmetry, *J. Phys. A: Math. Gen.*, 1999, **32**, pp.8785–8791, <http://xxx.lanl.gov/abs/cond-mat/9904360>
- [8] V.A.Avetisov, A.H.Bikulov, S.V.Kozyrev, V.A.Osipov, p -Adic Models of Ultrametric Diffusion Constrained by Hierarchical Energy Landscapes, *J. Phys. A: Math. Gen.*, 2002, **35**, pp.177–189, <http://xxx.lanl.gov/abs/cond-mat/0106506>

- [9] V.A.Avetisov, A.Kh.Bikulov, V.A.Osipov, p -Adic description of characteristic relaxation in complex systems, submitted to *J.Phys.A*, <http://xxx.lanl.gov/abs/cond-mat/0210447>
- [10] I.V.Volovich, p -Adic String, *Class. Quantum Gravity*, 1987, **4**, L83–L87
- [11] A.Khrennikov, p -Adic valued distributions in mathematical physics, Dordrecht, Kluwer Academic Publ., 1994
- [12] A.Khrennikov, *Non-Archimedean Analysis: Quantum Paradoxes, Dynamical Systems and Biological Models*, Dordrecht, Kluwer Academic Publishers, 1997
- [13] G.S.Djordjevic, B.Dragovich, *Mod.Phys.Lett.A*, 1997, **12**, p.1455
- [14] G.Parisi, N.Sourlas, p -Adic numbers and replica symmetry breaking, *European Phys. J. B*, 2000, **14**, p.535, <http://xxx.lanl.gov/abs/cond-mat/9906095>
- [15] D.M.Carlucci, C.De Dominicis, 1997, <http://xxx.lanl.gov/abs/cond-mat/9709200>
- [16] C.De Dominicis, D.M.Carlucci, T.Temesvari, *J. Phys. I France*, 1997, **7**, p.105
C.De Dominicis, D.M.Carlucci, T.Temesvari, 1997, <http://xxx.lanl.gov/abs/cond-mat/9703132>
- [17] M.Mezard, G.Parisi, N.Sourlas, G.Toulouse, M.Virasoro 1984 *Phys. Rev. Lett.* **52** 1156
- [18] K.Binder, A.P.Young 1986 *Rev. Mod. Phys.* **58** 801
- [19] M.Mezard, G.Parisi, M.Virasoro, *Spin-Glass Theory and Beyond*, Singapore, World Scientific, 1987
- [20] A.Ansary, J.Berendzen, S.F.Bowne, H.Frauenfelder, I.E.T.Iben, T.B.Sauke, E.Shyamsunder and R.D.Young 1985 *Proc. Natl. Acad. Sci. USA* **82** 5000
- [21] H.Frauenfelder, S.G.Sligar, P.G.Wolynes 1991 *Science* **254** 1598
- [22] H.Frauenfelder 1995 *Nature Struct. Biol.* **2** 821
- [23] D.Th.Leeson, D.A.Wiersma 1995 *Nature Struct. Biol.* **2** 848
- [24] D.Sherrington 1997 *Physica D* **107** 117
- [25] H.Frauenfelder, D.Th.Leeson 1998 *Nature Struct. Biol.* **5** 757
- [26] O.M.Becker, M.Karplus 1997 *J. Chem. Phys.* **106** 1495
- [27] J.P.Serre, *Trees*, New York, Berlin, Springer Verlag, 1980
- [28] A.T.Ogielski, D.L.Stein, *Phys.Rev.Lett.*, 1985, **55** N15, pp.1634-1637
- [29] Hajime Yoshino, Hierarchical diffusion, aging and multifractality, 1996, <http://xxx.lanl.gov/abs/cond-mat/9604033>